

Identification of Six Nassarid Snails Using *COI*-Based Restriction Fragment Length Polymorphism

Author(s): Yi Yang, Qi Li and Lingfeng Kong

Source: Journal of Shellfish Research, 37(1):239-243.

Published By: National Shellfisheries Association

URL: <http://www.bioone.org/doi/full/10.2983/035.037.0122>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

IDENTIFICATION OF SIX NASSARID SNAILS USING *COI*-BASED RESTRICTION FRAGMENT LENGTH POLYMORPHISM

YI YANG,¹ QI LI^{1,2*} AND LINGFENG KONG¹

¹Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, 5 Yushan Road, Qingdao 266003, China; ²Laboratory for Marine Fisheries Science and Food Production Processes, Qingdao National Laboratory for Marine Science and Technology, 1 Wenhai Road, Aoshanwei Town, Qingdao 266003, China

ABSTRACT The marine gastropod Nassariidae is a group of edible mud snails that are distributed worldwide. Some of the nassarid species could accumulate algal toxins that cause paralytic shellfish poisoning when being consumed. Therefore, efficient classification of nassarids is important for seafood safety, yet quite challenging because of their complex morphological diversity. In this study, an approach based on cytochrome c oxidase subunit 1–restriction fragment length polymorphism was developed to discriminate six common nassarid species, including *Nassarius nodifer*, *Nassarius conoidalis*, *Nassarius sinarus*, *Nassarius succinctus*, *Nassarius variciferus*, and *Reticunassa festiva*. Specific 709-bp fragments were polymerase chain reaction–amplified, which yielded differential digestion patterns after *MspI* restriction, allowing unambiguous discrimination of all six targeted species. Compared with a previously published approach for determination of nassarids using DNA barcodes, the approach developed here is much more efficient and cost saving, and will be an effective way to detect paralytic shellfish toxins–accumulating nassarids in seafood market.

KEY WORDS: identification, Nassariidae, paralytic shellfish poisoning, RFLP

INTRODUCTION

The mud snail Nassariidae are distributed worldwide on soft bottoms, with most species inhabiting intertidal regions from 0 to 300 m. The species diversity of nassarids is the highest in tropical regions, particularly the Indo-Pacific region (Cernohorsky 1972). In Asia, nassarids are consumed as seafood. Some nassarid species have been reported to contain algal toxins, especially paralytic shellfish toxins (Li & Chen 1981, Lin et al. 1998, Choi et al. 2006) and amnesic shellfish toxins (Goldberg 2003). These nassarid species have been widely recognized as a vector for paralytic shellfish poisoning (PSP) (Basti et al. 2018). Thus, consumption of such nassarids often leads to poisoning incidents in Asian countries, especially in China (Qin & Yu 2003, Shui et al. 2003, Yu et al. 2007, Luo et al. 2008, Zhang et al. 2009). Therefore, proper identification of such poisonous nassarids is necessary.

Identification of nassarid species is mainly based on shell morphology, especially the sculpture (Haasl 2000). The use of morphological characters for nassarid identification, however, is sometimes difficult because of intraspecific variation (Li et al. 2010). For example, the eurythermic species *Nassarius variciferus* (Adams, 1851) showed high intraspecific morphological diversity in the presence or absence of varices on body whorl and the color of spiral bands (Zou et al. 2012). Furthermore, interspecific convergence may also pose difficulties in species identification (Bargues & Mas-Coma 1997).

Molecular methods have been widely developed for species identification in the past two decades. The use of DNA-based techniques can overcome the drawbacks of morphological identification to a large extent (Tautz et al. 2003). Some examples of such methods include randomly amplified polymorphic DNA (Bardakci & Skibinski 1994), amplified fragment

length polymorphism (Congiu et al. 2002), DNA barcoding (Hebert et al. 2003a), and polymerase chain reaction (PCR)–based restriction fragment length polymorphism (RFLP) (Wolf et al. 1999).

Among these methods, PCR-RFLP relies on analysis of DNA fragment patterns after treatments with restriction enzymes and, therefore, is easy, rapid, and cost-effective. This method has been widely used for meat and fish identification, commonly using conserved mitochondrial (mt) DNA. Mitochondrial DNA presents a higher copy number and faster rate of mutation, making it generally more appropriate in the study for species identification (Chow et al. 2003). The most common mt genes used for species identification include *COI* (mt *COI*) (Haider et al. 2012, Mueller et al. 2015, Pappalardo & Ferrito 2015), cytochrome b (Dooley et al. 2005, Pepe et al. 2005), 12S ribosomal RNA (Fajardo et al. 2009), and 16S ribosomal RNA (Sumathi et al. 2015).

In the present study, a PCR-RFLP method using universal *COI* primers was developed for the identification of six common nassarid species, including *Nassarius nodifer*, *Nassarius conoidalis*, *Nassarius sinarus*, *Nassarius succinctus*, *Nassarius variciferus*, and *Reticunassa festiva*, along the coasts of China.

MATERIALS AND METHODS

Sample Collection and DNA Extraction

A total of 36 samples of *Nassarius nodifer*, *Nassarius conoidalis*, *Nassarius sinarus*, *Nassarius succinctus*, *Nassarius variciferus*, and *Reticunassa festiva* were collected along the coasts of China, with six individuals for each species. The species were first identified morphologically according to Cernohorsky (1972) and then confirmed by a molecular method (PCR-RFLP of *COI* sequences). Of the samples, the six *N. nodifer* were collected from a local market in

*Corresponding author. E-mail: qili66@ouc.edu.cn
DOI: 10.2983/035.037.0122

Zhanjiang (Guangdong Province of China), six *N. conoidalis* were collected from a local market in Dongshan County (Fujian Province of China), and six *N. sinarus* were collected from a local market in Zhoushan (Zhejiang Province of China). Other species, including *N. variciferus*, *N. succinctus*, and *R. festiva* (two individuals from each location), were collected from Binzhou (Shandong Province), Qingdao (Shandong Province), and Lianyungang (Jiangsu Province of China).

Specimens were stored in 95% ethanol. Genomic DNA was extracted from small pieces of foot tissue by the CTAB method as modified by Winnepenninckx et al. (1993).

Amplification of mt COI Gene Fragment

Universal *COI* primers (LCO1490: 5'-ATT CAA CCA ATC ATA AAG ATA TTG G-3' and HC02198: 5'-TAA ACT TCT GGA TGT CCA AAA AAT CA-3') were used for PCR amplification of all samples (Folmer et al. 1994). The standard PCRs were performed in a total volume of 25 μ L with 0.5 μ L of template DNA (approximately 100 ng), 2.5 μ L of 10 \times PCR buffer (Mg²⁺ plus), 0.5 μ L of dNTPs (10 mM), 1 μ L of each primer (10 μ M), and 0.25 μ L (5 U/ μ L) of *Taq* DNA polymerase. The following profile was used: predenaturation at 94°C for 3 min followed by 35 cycles of 94°C for 30 sec, 52°C annealing temperature for 30 sec, extension at 72°C for 1 min, and a final extension step at 72°C for 5 min. Generated PCR products were checked by electrophoresis on 1.5% agarose gel with ethidium bromide in 0.5 \times Tris Borate EDTA and were photographed under ultraviolet light using the gel documentation system.

DNA Sequencing and Selection of Restriction Enzymes

After PCR amplification, each product of *Nassarius nodifer*, *Nassarius conoidalis*, *Nassarius sinarus*, *Nassarius succinctus*, *Nassarius variciferus*, and *Reticunassa festiva* was sequenced by BGI sequencing, Beijing, China. These sequences were completed by retrieval from the corresponding mt genome, which were conducted in a previous analysis (unpublished) and submitted to GenBank with accession numbers KY783915–KY783920. The restriction enzymes *MspI* and *TaqI* were selected for the sequenced nucleotide using software Primer Premier 5 (Lalitha 2000). Gene alignment, recognition sites of two endonuclease, and expected product sizes are shown in Figure 1 and Table 1.

Polymerase Chain Reaction-Restriction Fragment Length Polymorphism Analysis

Digestion was performed with restriction enzymes *MspI* and *TaqI*. Digestion reactions were carried out in a total volume of 20 μ L containing 5 μ L of PCR product, 1 μ L of enzyme (10 units/ μ L), 2 μ L of 0.1% BSA, and 2 μ L of 10 \times *TaqI* buffer for *TaqI* or 2 μ L of 10 \times T buffer for *MspI*, respectively. The final volume was made up to 20 μ L with autoclaved sterile dH₂O. Restriction reactions were conducted in an incubator for 4 h at either 65°C for *TaqI* or 37°C for *MspI*. Digested products were run on 1.5% agarose gels and electrophoresed with 1 \times Tris acetate EDTA at 90 V for 45 min and visualized under ultraviolet light. A 100-bp DNA ladder was used as the reference marker.

RESULTS

A 709-bp fragment was amplified using universal *COI* primers and no size variations were found among different samples. For the analysis of digested products, only major bands were considered. The results showed that *TaqI* restricted positively in all samples (Fig. 2A, B), but failed to identify *Nassarius succinctus*, *Nassarius sinarus*, and *Nassarius conoidalis* because of the same length of restriction products. With respect to *MspI*, although it failed to restrict for *Reticunassa festiva*, all six nassarids exhibited distinctive restriction products (Fig. 2C, D). The length of the restriction products was in accordance with the size expected from the nucleotide sequences (Table 1).

DISCUSSION

Using only *Msp I* was sufficient to generate species-specific digested products to reveal levels of *COI* variation. According to Mueller et al. (2015), a mutation which affected an enzyme recognition site would lead to variations within samples of the same species; in the present analysis, no polymorphisms were detected when either *Taq I* or *Msp I* was applied to multiple samples of each species. This could be because of the conservation of mt *COI* gene within each species.

The DNA barcoding method has been proven as a useful tool for species identification in many animal and plant groups (Hebert et al. 2003a, 2003b). As for nassarid gastropods, Zou et al. (2012) reported a barcoding analysis by integrating mt and nuclear sequences and morphological characters, and successfully identified the species under study. Although this DNA sequencing-dependent technique was quite accurate and reliable, it was slow, costly, and unavailable for routine food species identification (Girish et al. 2005). Compared with DNA barcodes, PCR-RFLP was much more economical, easier, and faster and did not require high-quality or a large amount of DNA (Haider 2003). In addition, the materials and equipment, such as restriction enzymes, water bath, and electrophoresis system, were quite accessible for most biotechnology laboratories.

Previous studies using the PCR-RFLP method have been reported on the identification of livestock (Girish et al. 2005, Haider et al. 2012), fish (Mueller et al. 2015, Ferrito et al. 2016), microorganisms (Guillamón et al. 1998), and shellfish (Fernández-Tajes et al. 2011). Among these studies, many were related to fraudulent substitutions or adulterations of livestock meat or fish products. The present study focused on the identification of marine nassarids that may contain paralytic shellfish toxins. The toxins-containing species of nassarid in the China Sea have been discussed previously (Shui et al. 2003, Yu et al. 2007, Zhang et al. 2009). According to their results and records, most of the poisoning incidents via the consumption of nassarids were related to *Nassarius succinctus*, *Nassarius sinarus*, *Nassarius conoidalis*, and *Nassarius nodifer*. With respect to *Nassarius variciferus* and *Reticunassa festiva*, nonpoisoning accidents were found or recorded in these two species, and the widespread *N. variciferus* is still accessible in most seafood markets in China. In mainland China, about 30 species of nassarids have been reported (Yang & Zhang 2009). In the future, with more DNA data being sequenced, this method may be applied in the identification of PSP- versus non-PSP-accumulating species of nassarids.

<i>N. conoidalis</i>	TTTCTACAATCATAAAGATATTGGAACATTATATATCTATTTTGGAATATGATCAGGATTAGTTGGTACAGCTTTGAGGCTCTTATTCGTGCTGAAC	100
<i>N. nodifer</i>	TTTCTACAATCATAAAGACATTGGAACATTATATATCTATTTTGGAATATGATCAGGATTAGTTGGTACAGCTTTAAGACTCCTTATTCGTGCGGAAC	100
<i>N. sinarus</i>	TTTCTACAATCATAAAGATATTGGAACACTATATATCTATTTTGGAATATGATCAGGACTAGTTGGTACAGCCTTGAGACTTCTTATTCGTGCTGAAC	100
<i>N. succinctus</i>	TTTCTACAATCATAAAGACATTGGAACATTATATATCTATTTTGGAATATGATCAGGATTAGTTGGTACAGCCTTGAGACTTCTTATTCGTGCTGAAC	100
<i>N. variciferus</i>	TTTCTACAACCATAAAGACATTGGAACACTGTATATCTTTTCGGTATATGATCCGGATTAGTTGGGACAGCCTTAAGACTTCTTATTCGTGCTGAAT	100
<i>R. festiva</i>	TCCTACAATCATAAAGACATTGGAACGTTATATATTTTATTTGGAATATGATCAGGACTAGTTGGGACAGCTTTAAGCTTTTAACTTCGTGCTGAGTT	100
Consensus	t tctacaac cataaaga attggaac t tatatt t tt gg atatgac gga tagttgg ac gc tt ag ct t attcgtgc ga t	
<i>N. conoidalis</i>	TGGACAACCGGAGCTCTTCTGGTGACGATCAACTTTATAACGTAATTGTTACAGCTCATGCTTTTGAATAAATTTTTTCTAGTAATACCTATGATA	200
<i>N. nodifer</i>	TGGACAACCTGGAGCTCTTCTGGTGACGATCAACTTTATAACGTAATTGTTACAGCTCATGCTTTTGAATAAATTTTTTCTAGTAATACCTATGATA	200
<i>N. sinarus</i>	TGGGACGCTGGGACCTTCTGGGAGATGACCAACTTTATAATGTAATTGTTACGGCTCATGCTTTTGAATAAATTTTTCTTTAGTAATACCTATAATG	200
<i>N. succinctus</i>	TGGGCAACTGGAGCGCTTCTGGTGACGATCAACTTTATAACGTAATTGTTACAGCTCATGCTTTTGAATAAATTTTTTCTAGTAATACCTATGATA	200
<i>N. variciferus</i>	AGGCCAACCCAGGAGCACTCTTGGTGACGACCAACTTTATAATGTAATTGTTACAGCTCAGCCTTTTGAATAAATTTTTTCTTAGTAATACCTATAATG	200
<i>R. festiva</i>	AGGTCAACCTGGAGCACTTTTAGTGATGACCACTTTATAATGTAATTGTTACAGCTCATGCTTTTGAATAAATTTTTCTTTAGTAATACCTATGATA	200
Consensus	gg ca cc gg gc ctt t gg ga ga ca cttataa gtaattgttac gctca gcttttgaat atttt tt tagtaat cctat at	
<i>N. conoidalis</i>	ATTGGTGGATTGGGAATTGATTGGTTCGGCTAATATTAGGAGCTCCTGATATGGCTTTTCTCGTTAAATAATATAAGATTTTGATTACTTCTCTCCG	300
<i>N. nodifer</i>	ATTGGTGGATTTCGGAATTGATTAGTTCCGCTAATATTAGGAGCTCCTGATATGGCTTTTCTCGTTAAATAATATAAGATTTTGATTACTTCTCTCCG	300
<i>N. sinarus</i>	ATTGGTGGGTTTGGAAATGATTAGTTCCATTAATATTAGGAGCTCCTGATATAGCTTTTCTCGTTAAATAATATAAGATTTTGATTACTTCTCTCCG	300
<i>N. succinctus</i>	ATTGGTGGATTGGGAATTGATTAGTTCCGCTAATATTAGGAGCTCCTGATATAGCTTTTCTCGTTAAATAATATAAGATTTTGATTACTTCTCTCCG	300
<i>N. variciferus</i>	ATTGGGGATTGGGAATTGATTAGTTCCCTCTTATGTTAGGAGCTCCTGATATGGCTTTTCTCGATTAAATAATATAAGTTTGGATTACTTCTCTCCG	300
<i>R. festiva</i>	ATTGGAGGTTTGGAAATGGTTAGTTCCGTTGATATTGGGAGCTCCTGATATAGCTTTTCTCGACTAAATAACATAAGATTTTGATTGCTTCTCTCCG	300
Consensus	attgg gg tt gg aattg tt gtcc t at tt ggaactcc gatat gcttt cctcg taaataa ataag ttttg tt cttctcc g	
<i>N. conoidalis</i>	CACTACTCTTTTATTATCTTCTGCCGAGTAGAAAAGGGGTAGGAACCGGATGAACGGTTTATCCCTTTATCTGCTAATTTAGCTCATGCTGGTGG	400
<i>N. nodifer</i>	CATTACTCTTTTATTATCTCTGCCGAGTAGAAAAGGGGTAGGAACCTGGATGGACTGTTTATCCCTTTATCTGCTAATTTAGCTCATGCTGGTGG	400
<i>N. sinarus</i>	CATTACTCTTTTATTATCTTCTGCCAGCAGTAGAAAAGGGGTAGGAACCTGGATGAACGGTTTATCCCTTTATCTGCTAATTTAGCTCATGCTGGTGG	400
<i>N. succinctus</i>	CATTACTCTTTTGGTTATCTTCTGCCAGCAGTAGAAAAGGGGTAGGAACCTGGATGAACCTGTTTATCCCTTTATCTGCTAATTTAGCTCATGCTGGTGG	400
<i>N. variciferus</i>	CTTTACTCTTTTACTATCTTCCAGCTGCTGTTGAAAGGGGGTGGGACTGGTGAACCTGTTTATCCCTTTATCTGCTAATTTAGCTCATGCTGGAGG	400
<i>R. festiva</i>	CACCTTTACTTTTACTATCTTCCAGCAGCAGTAGAAAAGGGGGTACTGGATGAACCTGTTTATCCCTTTATCTGCTAATTTAGCTCATGCTGGTGG	400
Consensus	c t t ctttt tadc tc gc gc gt gaaag gg g gg ac gg tg ac gt ta cc cttttatct taat t gctca gc gg gg	
<i>N. conoidalis</i>	TTCTGTAGATTTAGCTATTTTTCTTTACATCTTGCAGGTGATCATCAATTCTTGGGCTGTTAATTTTATACGACTATTATAATATACGATGACGA	500
<i>N. nodifer</i>	TTCTGTAGATTTAGCTATTTTTCTTTACATCTTGCAGGTGATCATCAATTCTTGGGCTGTTAATTTTATACGACTATTATAACATGCGATGACGA	500
<i>N. sinarus</i>	TTCTGTAGATTTAGCCATTTTTCTTTACATCTTGCAGGTGATCATCAATTCTTGGAGCTGTTAATTTTATACGACTATTATAATATACGATGACGG	500
<i>N. succinctus</i>	TTCTGTGATTTAGCTATTTTTCTTTACATCTTGCAGGTGATCATCAATTCTTGGGCTGTTAATTTTATACACTATTATAATATACGATGACGA	500
<i>N. variciferus</i>	TTGAGTATTTAGCTATTTTTCTTCTTACATCTTGCAGGTGTTTCTTCTATTTTAGGAGCTGTAACCTTTATACAACCTATTATAACATACGATGACGT	500
<i>R. festiva</i>	TTGAGTTGACTTAGCAATTTTTCACTTCAATTTAGCAGGTGATCTTCGAATTTTAGGGCTGTAACCTTTATACAACCTATTATAATATACGATGACGA	500
Consensus	ttc gt ga ttacg attttttc t cat t gc ggtgt tc ac att t gg gctgt aa tttat ac actattattaa at cgatg cg	
<i>N. conoidalis</i>	GGAAATCAGTTTGAACGCTCTCCTTTATTTGATGATCTGTTAAAATTACTGCTATTTTACTTCTTTTATCTTTACCAGTGTAGCAGGAGCTATTACAA	600
<i>N. nodifer</i>	GGGATGCAATTCGAACGCTCTCCTTTATTTGATGATCTGTTAAAATTACTGCTATTTTACTTCTTTTATCTTTACCAGTATTAGCAGGAGCTATTACAA	600
<i>N. sinarus</i>	GGAAATGCAATTTGAACGCTCTCCTTTATTTGATGATCTGTTAAAATTACTGCTATTTTACTTCTTTTATCTTTACCAGTATTAGCAGGAGCTATTACAA	600
<i>N. succinctus</i>	GGAAATGCAATTTGAACGCTCTCCTTTATTTGATGATCTGTTAAAATTACTGCTATTTTACTTCTTTTATCTTTACCAGTATTAGCAGGAGCTATTACAA	600
<i>N. variciferus</i>	GGAAATGCAATTTGAGCGCTCTCCTTTATTTGATGATCTGTTAAAATTACTGCTATTTTACTTCTTTTATCTTTACCAGTATTAGCAGGAGCTATTACAA	600
<i>R. festiva</i>	GGGATCAGTTTGAACGCTCTCCTTTATTTGATGATCTGTTAAAATTACTGCTATTTTACTTCTTTTATCTTTACCAGTATTAGCAGGAGCTATTACAA	600
Consensus	gg at ca tt ga cg ct cct tatttgt tgatc gt aaaat actgctatttta t ctt t tct t cc gt t gc ggagctat acaa	
<i>N. conoidalis</i>	TGCTTCTAAGTGAACGAAATTTAATACTGCGTCTTTGATCCAGCAGGTGGTGGAGATCCTATTTTATATCAACATTTGTTTGTATTTTGGGCATCC	700
<i>N. nodifer</i>	TACTTTTAACTGAACGAAATTTAATACTGCATTTTGTATCCGCGAGGAGTGGAGATCCTATTTTATATCAACATTTATTTGATTTTGGGCATCC	700
<i>N. sinarus</i>	TATTATTAAGTGAACGAAATTTAATACTGCATTTTGTATCCGCGAGGAGTGGAGATCCTATTTTATATCAACATTTATTTGATTTTGGGCATCC	700
<i>N. succinctus</i>	TATTATTAAGTGAACGAAATTTAATACTGCATTTTGTATCCAGCTGGAGGTGGAGATCCTATTTTATACCAACATTTATTTGATTTTGGGCATCC	700
<i>N. variciferus</i>	TACTGTTAAGTGAACGAAATTTAATACTGCATTTTGTATCCGCGAGGAGTGGAGATCCTATTTTATATCAACATTTATTTGGTTTTGGGCATCC	700
<i>R. festiva</i>	TACTTCTAAGCAGCAAAATTTAATACTGCTTTTTCGAATTTTAGCAGGAGGTTGGTATCCAAATTTTATACCAACATTTATTTGATTTTGGGCATCC	700
Consensus	t t taac ga cgaaattttaa ac gc tt tt ga cc gc gg ggtgg gatcc attttata caaca t tttg tt ttgg ca cc	
<i>N. conoidalis</i>	TGAAGTTTA	709
<i>N. nodifer</i>	TGAAGTTTA	709
<i>N. sinarus</i>	TGAAGTTTA	709
<i>N. succinctus</i>	TGAAGTTTA	709
<i>N. variciferus</i>	TGAGGTATA	709
<i>R. festiva</i>	TGAAGTCTA	709
Consensus	tga gt ta	

Figure 1. Partial nucleotide sequences of the mt *COI* gene used for the RFLP analysis of six nassarid species. Recognition sites of *MspI* (C^{CGG}) are underlined, whereas those of *TaqI* (T^{CGA}) are in italics and bold style.

It can be concluded that the six nassarid species can be qualitatively identified and differentiated by the quick, easy, low-cost, and reliable PCR-RFLP of the mt *COI* gene.

This method could also be a potential way for future monitoring of PSP-accumulating nassarid species in the seafood market.

TABLE 1.

Expected size (bp) of digested DNA fragments on digestion with two restriction enzymes.

Enzyme	Species					
	<i>Reticunassa festiva</i>	<i>Nassarius variciferus</i>	<i>Nassarius succinctus</i>	<i>Nassarius nodifer</i>	<i>Nassarius sinarus</i>	<i>Nassarius conoidalis</i>
<i>MspI</i>	Negative	55, 63, 70, 521	315, 394	67, 642	63, 70, 576	109, 240, 360
<i>TaqI</i>	72, 184, 190, 263	95, 263, 351	95, 614	95, 103, 511	95, 614	95, 614




Figure 2. Restriction profiles of *COI* amplicons with two restriction enzymes in the nassariid samples analyzed. Lanes 8, 16, and 24, 100-bp DNA ladder. (A) *TaqI*. Lanes 1, 9, and 17, unrestricted PCR product of *Nassarius variciferus*, *Nassarius nodifer*, and *Reticunassa festiva*, respectively; lanes 2–7, 10–15, and 18–23, restriction profiles generated by *N. variciferus*, *N. nodifer*, and *R. festiva*, respectively. (B) *TaqI*. Lanes 1, 9, and 17, unrestricted PCR product of *Nassarius succinctus*, *Nassarius Conoidalis*, and *Nassarius sinarus*, respectively; lanes 2–7, 10–15, and 18–23, restriction profiles generated by *N. succinctus*, *N. conoidalis*, and *N. sinarus*, respectively. (C) *MspI*. Lanes 1, 9, and 17, unrestricted PCR product of *N. conoidalis*, *N. variciferus*, and *N. succinctus*, respectively; lanes 2–7, 10–15, and 18–23, restriction profiles generated by *N. conoidalis*, *N. variciferus*, and *N. succinctus*, respectively. (D) *MspI*. Lanes 1, 9, and 17, unrestricted PCR product of *R. festiva*, *N. sinarus*, and *N. nodifer*, respectively; lanes 2–7, 10–15, and 18–23, restriction profiles generated by *R. festiva*, *N. sinarus*, and *N. nodifer*, respectively.

ACKNOWLEDGMENTS

This study was supported by grants from the Doctoral Program of the Ministry of Education of China

(20130132110009), Fundamental Research Funds for the Central Universities, and Qingdao National Laboratory for Marine Science and Technology (2015ASKJ02).

LITERATURE CITED

- Bardakci, F. & D. O. F. Skibinski. 1994. Application of the RAPD technique in tilapia fish: species and subspecies identification. *Heredity* 73:117–123.
- Bargues, M. D. & S. Mas-Coma. 1997. Phylogenetic analysis of Lymnaeid snails based on 18S rDNA sequences. *Mol. Biol. Evol.* 14:569–577.
- Basti, L., H. Hégaret & S. E. Shumway. 2018. Harmful algal blooms and shellfish. In: Shumway, S. E., J. M. Burkholder & S. Morton, editors. *Harmful Algal Blooms: A Compendium Desk Reference*. UK: John Wiley & Sons.
- Cernohorsky, W. O. 1972. Indo-Pacific Nassariidae (Mollusca: Gastropoda). *Rec. Auckland Mus.* 9:125–194.
- Choi, M. C., K. N. Peter, D. P. Hsieh & P. K. Lam. 2006. Trophic transfer of paralytic shellfish toxins from clams (*Ruditapes philippinarum*) to gastropods (*Nassarius festivus*). *Chemosphere* 64:1642–1649.
- Chow, S., T. Nohara, T. Tanabe, S. Itoh, T. Tsuji & Y. Nihikawa. 2003. Genetic acid morphological identification of larvae and small juvenile tunas (*Pisces scombridae*) caught by mid-water trawl in the eastern Pacific. *Bull. Fish. Res. Agen.* 8:1–14.
- Congiu, L., F. Fontana, T. Patarnello, R. Rossi & L. Zane. 2002. The use of AFLP in sturgeon identification. *J. Appl. Ichthyology* 18:286–289.
- Dooley, J. J., H. D. Sage, M. A. L. Clarke, H. M. Brown & S. D. Garrett. 2005. Fish species identification using PCR-RFLP analysis and lab-on-a-chip capillary electrophoresis: application to detect white fish species in food products and an interlaboratory study. *J. Agric. Food Chem.* 53:3348–3357.
- Fajardo, V., I. González, J. Dooley, S. Garret, H. M. Brown, T. García & R. Martín. 2009. Application of polymerase chain reaction-restriction fragment length polymorphism analysis and lab-on-a-chip capillary electrophoresis for the specific identification of game and domestic meats. *J. Sci. Food Agric.* 89:843–847.
- Fernández-Tajes, J., A. Longa, J. García-Gil, Y. W. Chiu, Y. S. Huang, J. Méndez & R. S. Lee. 2011. Alternative PCR-RFLP methods for mussel *Mytilus* species identification. *Eur. Food Res. Technol.* 233:791.
- Ferrito, V., V. Bertolino & A. M. Pappalardo. 2016. White fish authentication by COI-Bar-RFLP: toward a common strategy for the rapid identification of species in convenience seafood. *Food Control* 70:130–137.
- Folmer, O., M. Black, W. Hoeh, R. Lutz & R. Vrijenhoek. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* 3:294–299.
- Girish, P. S., A. S. R. Anjaneyulu, K. N. Viswas, B. M. Shivakumar, M. Anand, M. Patel & B. Sharma. 2005. Meat species identification by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) of mitochondrial 12S rRNA gene. *Meat Sci.* 70:107–112.
- Goldberg, J. D. 2003. Domoic acid in the benthic food web. Master's Thesis, California State University, Monterey Bay, CA. 33 pp.
- Guillamón, J. M., J. Sabaté, E. Barrio, J. Cano & A. Querol. 1998. Rapid identification of wine yeast species based on RFLP analysis of the ribosomal internal transcribed spacer (ITS) region. *Arch. Microbiol.* 169:387–392.
- Haasl, D. M. 2000. Phylogenetic relationships among Nassariid gastropoda. *J. Paleontol.* 74:839–852.
- Haider, N. 2003. Development and use of universal primers in plants. PhD thesis, University of Reading, UK.
- Haider, N., I. Nabulsi & B. Al-Safadi. 2012. Identification of meat species by PCR-RFLP of the mitochondrial COI gene. *Meat Sci.* 90:490–493.
- Hebert, P. D. N., A. Cywinska, S. L. Ball & J. R. deWaard. 2003a. Biological identifications through DNA barcodes. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* 270:313–321.
- Hebert, P. D. N., S. Ratnasingham & J. R. deWaard. 2003b. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* 270: S96–S99.
- Lalitha, S. 2000. Primer premier 5. *Biotech Softw. Internet Rep.* 1:270–272.
- Li, H., D. Lin, H. Fang, A. Zhu & Y. Gao. 2010. Species identification and phylogenetic analysis of genus *Nassarius* (Nassariidae) based on mitochondrial genes. *Chin. J. Oceanology Limnol.* 28:565–572.
- Li, Y. & S. Chen. 1981. Report of 8 incidents of intoxication due to *Nassarius* (*Zeuxis*) *succinctus*. *Chinese J. Surg.* 20:231–232 (in Chinese).
- Lin, Y., X. Jia, M. Yang, Y. Zhong, G. Quan & P. Fan. 1998. Paralytic shellfish poison in contaminated shellfish along coast of China. *Tropic Oceanology* 18:90–96.
- Luo, X., R. C. Yu, X. J. Wang, J. Li & M. J. Zhou. 2008. Toxicity dynamics and toxin composition of *Nassarius* spp. from Fujian province. *J. Fish. Chin.* 32:636–643.
- Mueller, S., S. M. Handy, J. R. Deeds, G. O. George, W. J. Broadhead, S. E. Pugh & S. D. Garrett. 2015. Development of a COX1 based PCR-RFLP method for fish species identification. *Food Control* 55:39–42.
- Pappalardo, A. M. & V. A. Ferrito. 2015. COI-Bar-RFLP strategy for the rapid detection of *Engraulis encrasicolus* in processed anchovy products. *Food Control* 57:385–392.
- Pepe, T., M. Trotta, I. Di Marco, P. Cennamo, A. Anastasio & M. L. Cortesi. 2005. Mitochondrial cytochrome b DNA sequence variations: an approach to fish species identification in processed fish products. *J. Food Prot.* 68:421–425.
- Qin, P. Z. & M. Yu. 2003. Research on Nassidae food poisoning in Ningbo City from 1985 to 2000. *Chin. J. Prev. Med.* 37:226.
- Shui, L. M., K. Chen, J. Y. Wang, H. Z. Mei, A. Z. Wang, Y. H. Lu & D. F. Hwang. 2003. Tetrodotoxin-associated snail poisoning in Zhoushan: a 25-year retrospective analysis. *J. Food Prot.* 66:110–114.
- Sumathi, G., G. Jeyasekaran, R. J. Shakila, B. Sivaraman, G. Arunkumar, U. Manimaran & D. Sukumar. 2015. Molecular identification of grouper species using PCR-RFLP technique. *Food Control* 51:300–306.
- Tautz, D., P. Arctander, A. Minelli, R. H. Thomas & A. P. Vogler. 2003. A plea for DNA taxonomy. *Trends Ecol. Evol.* 18:70–74.
- Winnepeninckx, B., T. Backeljau & R. Dewachter. 1993. Extraction of high molecular weight DNA from molluscs. *Trends Genet.* 9:407.
- Wolf, C., J. Rentsch & P. Hübner. 1999. PCR-RFLP analysis of mitochondrial DNA: a reliable method for species identification. *J. Agric. Food Chem.* 47:1350–1355.
- Yang, J. W. & S. P. Zhang. 2009. Status and prospects of the systematics of Nassariidae (Mollusca: Gastropoda) in China coasts. *Mark. Sci.* 33:123–126.
- Yu, R. C., M. J. Zhou, A. F. Li, Q. C. Zhang, Y. F. Wang, J. Li, M. Quilliam & B. Luckas. 2007. Analysis of toxins in snails responsible for two poisoning incidents in China. *J. Fish. Sci. Chin.* 14:801–806.
- Zhang, N., J. Su, H. X. Liu, S. Z. Ye, L. F. Li & L. Z. Cai. 2009. The species and toxicities of Nassariidae collected from the coast of Southeast China Sea. *Asian J. Ecotox.* 4:289–294.
- Zou, S. M., Q. Li & L. F. Kong. 2012. Monophyly, distance and character-based multigene barcoding reveal extraordinary cryptic diversity in *Nassarius*: a complex and dangerous community. *PLoS One* 7:e47276.